

## Abstract


### Background

Autotaxin (ATX) drives stromal remodeling, fibrosis, and immune suppression in solid tumours. Cambritaxestat (IOA-289) is an oral, selective ATX inhibitor. An AI-driven reverse translational framework prioritised hepatocellular carcinoma (HCC) as a leading indication for ATX targeting. We further evaluated ATX expression and cambritaxestat activity in human HCC and cholangiocarcinoma (CC) using complementary translational platforms benchmarked against standard of care (SoC).

### Results

AI analysis identified HCC as the top candidate indication. ATX expression was quantified in HCC tumouroids, and five ATX-high models were selected for tumour cell killing assays. Cambritaxestat was tested as monotherapy or combined with bevacizumab plus durvalumab (standard of care (SoC) and compared with SoC alone; tumour cell death was assessed by confocal imaging. In ATX-high HCC tumouroids, cambritaxestat monotherapy induced tumour cell death in 80% (4/5) of models, compared with 60% (3/5) for SoC. Combination therapy achieved responses in 80% (4/5) of models. In an independent platform, combination treatment was predicted to outperform SoC across all HCC samples (4/4), while cambritaxestat monotherapy reached response thresholds in 50% (2/4). In CC, combination therapy induced responses in 60% (3/5) versus 20% (1/5) in SoC alone.


## Cambritaxestat Mechanism of Action



## Conclusion


Ranking of indications: AI based search of online databases and clinical studies and by measuring ATX/LPAR expressions and LPA levels

Efficacy was shown in HCC tumouroids by measuring direct cell killing and immune system activation




Across independent human tumouroid platforms, cambritaxestat demonstrated antitumour activity as monotherapy in HCC and enhanced efficacy in combination with SoC in both HCC and CC. These findings support further clinical development of ATX inhibition in liver cancers.


## 1. AI-Driven approach identifies HCC as top indication candidate based on known literature




## 2. Public dataset and Biobanks analyses support a rationale to focus on liver indications




## 3. Optimized anti-ATX/ENPP2 IHC staining of human hepatocellular carcinoma tissue and Cholangiocarcinoma tissue



## 4. Cambritaxestat (IOA-298) mono- and combination therapy induce tumour cell death in human HCC tumouroids



## 5. Cambritaxestat is predicted to improve SoC efficacy in HCC and iCCA in human tumours



## 6. Two independent preclinical models performed in human HCC tumouroids show consistent cambritaxestat response in monotherapy or in combination with SoC

| A                   | Comparison HCC                                                                                 |                       |
|---------------------|------------------------------------------------------------------------------------------------|-----------------------|
|                     | Nilogen Oncosystems                                                                            | Calico Biosystems     |
| HCC:                | Sample selection by ATX expression: Yes, over 30% expression by IHC. No                        |                       |
|                     | Readout: Direct tumor cell killing → intrinsic. Immune microenvironment parameters → extrinsic |                       |
| IOA-289 Monotherapy | 4/5 responders (80%)                                                                           | 2/4 responders (50%)  |
|                     | IOA-289 + SoC                                                                                  | 4/4 responders (100%) |
| SoC                 | 3/5 responders (60%)                                                                           | 1/4 responders (25%)  |

**Acknowledgments:**  
The authors thank Riccardo Turrini and Marie-Agnes Doucey (Calico Biosystems), William Coward (Nilogen Oncosystem), and Nives Rombini and Jonas Walheim (Navis Bio), together with all study personnel from iOnctura BV and iOnctura SA. This study was sponsored by iOnctura BV.